

## Features

- 1 Incorrect administration is prevented by removing part of the label and attaching it to the syringe after drawing the medical solution from the vial.
- 2 Also, when co-infusing the medical solution into an infusion bag, part of the removed label can be attached and forgetting to perform infusion as well as double infusion can be prevented.
- 3 Due to the use of reusable adhesive, re-attachment is possible based on usage history from vial → syringe → medical chart.
- 4 Substantial equipment investment is not necessary due to the ability to use existing labelers in most cases.
- 5 The shape and position of the removal section can be created according to demand.

## Usage method



2

3

*To Prevent Medication Error*

# Pre-use adhesive label For vials



IWATA LABEL CO., LTD.



Head office /  
Nagoya office

5-15-18 Mitsui, Ichinomiya City, Aichi Prefecture 491-0827  
TEL (0586)76-1612 FAX (0586)76-1724

Tokyo office

(Asakusabashi Sugiura Building 4F) 1-23-4 Yanagibashi, Daito Ward, Tokyo 111-0052  
TEL (03)3862-9051 FAX (03)3862-9056

Osaka office

(Meiyu Building 5F) 1-22-22 Esaka-cho, Suita City, Osaka 564-0063  
TEL (06)6339-4401 FAX (06)6339-4403

Toyama office

(La Fontel Building 6F) 7-9 Uchisaiwai-cho, Toyama City, Toyama Prefecture 30-0093  
TEL (076)442-1997 FAX (076)442-1980

Akita office

(Castle Oak A-5) 3-2-48-8 Katayama-cho, Odate City, Akita Prefecture 017-0872  
TEL (0186)42-2608 FAX (0186)42-2608

<http://www.iwatalabel.co.jp>